Amendment to the Claims

Please amend claims 6-8 and 13-15 and add new claims 16-20 as shown below.

- 1. (Original) A method for inspecting fat-soluble vitamins and/or fat-soluble food factors in a body, comprising analyzing fat-soluble vitamins and/or fat-soluble food factors in saliva.
- 2. (Original) A method for inspecting migration of fat-soluble vitamins and/or fat-soluble food factors contained in ingested health supplements, drugs or foods into a body, comprising analyzing fat-soluble vitamins and/or fat-soluble food factors in saliva.
- 3. (Original) The method according to claim 2, wherein the inspecting is conducted on the basis of concentrations of the fat-soluble vitamins and/or the fat-soluble food factors in saliva of subjects before ingestion of the health supplements, drugs or foods, or the mean concentrations of the fat-soluble vitamins and/or the fat-soluble food factors in saliva of a control group ingesting no health supplements, drugs or foods.
- 4. (Original) A method for inspecting effect or action of an administered therapeutic agent on biosynthesis and metabolism of fat-soluble vitamins and/or fat-soluble food factors, comprising analyzing fat-soluble vitamins and/or fat-soluble food factors in saliva as indicators.
- 5. (Original) The method according to claim 4, wherein the inspecting is conducted on the basis of concentrations of the fat-soluble vitamins and/or the fat-soluble food factors in saliva of subjects before taking the therapeutic agent, or the mean concentrations of the fat-soluble vitamins and/or the fat-soluble food factors in saliva of a control group with no taken therapeutic agent.

- 6. (Currently Amended) A method for assessing suitability of ingestion or an intake of health supplements, drugs or foods comprising fat-soluble vitamins and/or fat-soluble food factors, wherein the method uses a method according to claim 2 or 3.
- 7. (Currently Amended) A method for assessing suitability of administration or a dosage of a therapeutic agent having effect or action on biosynthesis and metabolism of fat-soluble vitamins and/or the fat-soluble food factors, wherein the method uses a method according to claim 4 or 5.
- 8. (Currently Amended) The method according to any of claims claim 1 to 7, wherein saliva is parotid saliva.
- 9. (Original) The method according to claim 8, wherein the parotid saliva is collected using a saliva collecting tool which collects parotid saliva selectively and quantitatively.
- 10. (Original) The method according to claim 9, wherein the saliva collecting tool has (a) a collection part comprising an absorber for absorbing saliva irreversibly and (b) a quantification part for quantifying the amount of saliva collected to the absorber.
- 11. (Original) The method according to claim 10, wherein the saliva collecting tool has further a storage container part for holding a saliva preservative solution in which the absorber with the absorbed saliva is dipped to preserve.
- 12. (Original) The method according to claim 11, wherein the saliva preservative solution is a water-soluble organic solvent.
- (Currently Amended) The method according to any of claims claim 1 to 12, comprising:(a) a process for extracting the collected saliva, the absorber with the absorbed saliva,and/or the preservative solution with the preserved absorber with a water-soluble organic

- solvent, a hydrocarbon base organic solvent, or a mixture of isopropanol and ethyl acetate,
- (b) a process for using the extract as an analysis sample to separate the fat-soluble vitamins and/or the fat-soluble food factors by high-performance liquid chromatography, and
- (c) a process for detecting the separated fat-soluble vitamins and/or the fat-soluble food factors.
- 14. (Currently Amended) The method according to any of claims claim 1 to 13, wherein the fat-soluble vitamins and/or the fat-soluble food factors are at least one substance selected from a group consisting of CoQ10, lycopene, β-carotene and tocopherol.
- 15. (Currently Amended) A method for screening drugs or health supplements, wherein the method uses a method according to any of claims claim 1 to 14.
- 16. (New) A method for assessing suitability of ingestion or an intake of health supplements, drugs or foods comprising fat-soluble vitamins and/or fat-soluble food factors, wherein the method uses a method according to claim 3.
- 17. (New) A method for assessing suitability of administration or a dosage of a therapeutic agent having effect or action on biosynthesis and metabolism of fat-soluble vitamins and/or the fat-soluble food factors, wherein the method uses a method according to claim 5.
- 18. (New) The method according to claim 12, comprising:
 - (a) a process for extracting the collected saliva, the absorber with the absorbed saliva, and/or the preservative solution with the preserved absorber with a water-soluble organic

solvent, a hydrocarbon base organic solvent, or a mixture of isopropanol and ethyl acetate,

- (b) a process for using the extract as an analysis sample to separate the fat-soluble vitamins and/or the fat-soluble food factors by high-performance liquid chromatography, and
- (c) a process for detecting the separated fat-soluble vitamins and/or the fat-soluble food factors.
- 19. (New) The method according to claim 13, wherein the fat-soluble vitamins and/or the fat-soluble food factors are at least one substance selected from a group consisting of CoQ10, lycopene, β-carotene and tocopherol.
- 20. (New) A method for screening drugs or health supplements, wherein the method uses the method according to claim 14.